


Tree phenology in relation to meteorological conditions and crown defoliation on forest monitoring plots in Slovenia



Urša Vilhar, Mitja Skudnik, Mitja Ferlan, Primož Simončič



GOZDARSKI INŠTITUT SLOVENIJE
SLOVENIAN FORESTRY INSTITUTE





 **ICP Forests**

Introduction



- Linking tree phenology to crown defoliation and meteorological conditions
 - *Fagus sylvatica*, *Quercus robur* and *Picea abies*
 - 9 forest monitoring plots in Slovenia in 2004 – 2013

Published in: Vilhar U, M Skudnik, M Ferlan, P Simončič (2014) Influence of meteorological conditions and crown defoliation on tree phenology in intensive forest monitoring plots in Slovenia. *Acta Silvae et Ligni* 105:1-15





GOZDARSKI INŠTITUT SLOVENIJE
SLOVENIAN FORESTRY INSTITUTE



ICP Forests

Materials and methods

- Tree phenology
 - The first leaf and needle unfolding (LU)
 - The general leaf colouring (LC) and
 - The length of the growing season (LGS)
- Crown defoliation
 - the amount of foliage missing in comparison to a full-leaved reference tree, estimated in 5% classes for individual trees from 1-3 social class



Materials and methods




- Meteorological conditions
 - Monthly and three-monthly sums of precipitation (P)
 - Monthly means, monthly and three-monthly sums of air temperature (T)
- Relative extractable soil water
 - Monthly means of relative extractable water (REW)
 - Calculated from the daily minimum and maximum soil water content ($\text{m}^3 \text{m}^{-3}$)

$$REW = \frac{SWC_{day} - SWC_{min}}{SWC_{max} - SWC_{min}}$$




Results


- A high sensitivity of first leaf unfolding to air temperature and precipitation for all species, exhibiting contrasting responses



Needle and leaf unfolding (LU)



Autumn coloring of leaves (LC)





MANAGING FORESTS
FOR MULTIPLE PURPOSES:
CARBON BIODIVERSITY SOCIO-ECONOMIC WELLBEING



Results





- A strong sensitivity of beech defoliation to P and REW



Results




- Higher crown defoliation of pedunculate oak might contribute to earlier LU, later LC and longer LGS in the next year.



Conclusions




- Oak crown defoliation and next-year phenology were correlated, with higher crown defoliation contributing to earlier leaf unfolding, later autumn leaf coloring and longer growing season in next year.
- No correlation was found between phenology and crown defoliation for beech nor spruce.
- To assess the influence of crown defoliation and meteorological conditions on tree phenology, longer time series are needed, involving a larger number of tree species and sites.



Acknowledgments

- The study was part of:
 - ManFor C. BD project, titled “Managing forests for 649 multiple purposes: carbon, biodiversity and socio-economic wellbeing” (LIFE09ENV/IT/000078),
 - EUFORINNO - »European Forest Research and Innovation« (Reg.Pot No. 315982),
 - Program group Forest biology, ecology and technology, financed by Ministry of Education, Science and Sport of the Republik of Slovenia.



GOZDARSKI INŠTITUT SLOVENIJE
SLOVENIAN FORESTRY INSTITUTE

