

# Impact of N inputs on forest and forest soils biogeochemistry in Great Britain

Sue Benham



Elena Vanguelova & Rona Pitman

Centre for Forestry and Climate Change  
Forest Research, UK



## Main Concerns

- Critical N load exceeded in >70% of broadleaf woodland and conifer woodland in the UK,
- The importance of small scale N variability and its impact, e.g. at woodland edges due to the intensity of UK agriculture, farming, road traffic and the fragmented nature of the forests.
- Changes in nitrogen and carbon of forest and forest soils and the possibilities of  $\text{NO}_3$  leaching to waters, the implications for forest GHG mitigation potential and the impact on biodiversity.





## N inputs affecting forests and soils

- N deposition by local, regional and national sources (intensive farming, agriculture and intensive road system)
- Forest fertilisation (in the past but with long term effects on soil and forest production)
- Other N generated input (such as canopy N generated by biological agents)



## This overview includes:

- Small scale impacts of N deposition on soil and forest biochemistry under both broadleaf and conifer stands from N gradients studies (farming and roads)
- Regional differences in N impacts by comparing areas with low and high N deposition
- Long term monitoring of N inputs and outputs and its effects on the forest and soils from the intensive ICP forest monitoring network
- Forest soil N status from extensive national soil survey (BioSoil)

## Chicken and pig sheds in Thetford






## Experimental transects under Corsican pine and beech stands away from pig and chickens farms

### Measurements along transects:

- Air quality ( $\text{NH}_3$ )
- Rainfall and throughfall chemistry
- Litterfall biomass and chemistry
- Soil solution chemistry
- Litter decomposition
- Foliar chemistry
- Soil chemistry
- Tree growth

Averaged annual rainfall - 600mm  
Sandy soil over chalk






## Monitoring at Thetford







## Gradient and regional studies - at Thetford and Alice Holt forests



*Estimated distribution of dry deposition of ammonia to semi-natural vegetation.*


*Estimated distribution of wet deposition of ammonium in rain and snow.*

Thetford N deposition 18 to 47 kg/ha/a; Alice Holt 8 to 12 kg/ha/a






## Air NH<sub>3</sub> at Thetford and Alice Holt



## Total N deposition (throughfall) at Thetford and Alice Holt






Thetford N deposition 18 to 47 kg/ha/a; Alice Holt 8 to 12 kg/ha/a

Vanguelova and Pitman, in prep.






## Soil pH, NH<sub>4</sub>-N, NO<sub>3</sub>-N in Thetford's gradients

### Corsican pine




### Beech






## Foliar N in Corsican pine and beech at Thetford and Alice Holt forest

Corsican  
pine








Beech








## Litterfall biomass at Thetford and Alice Holt Corsican pine and Beech forests








## Litter decomposition under Corsican pine at Thetford





## Tree annual growth at Alice Holt and Thetford forests




Does N exacerbate the drought impact on tree growth?



Vanguelova and Pitman, in prep.





## Extensive and Intensive ICP forest monitoring networks






Observed  
relationships  
between forest  
condition and  
nitrogen  
deposition  
(Level I)

Kennedy and Freer- Smith, 2000





## Biological N inputs (Level II)

*Tortrix viridana*






*Operophtera brumata*



Pitman et al., 2010



## Changes in soil solution $\text{NO}_3$ and $\text{NH}_4$ in 9 Level II sites (1994-2006)




- Decrease in soil solution  $\text{NO}_3$  only at sites with N deposition or overall acidic pollution decline
- $\text{NO}_3$  concentrations in the soils are very dependent on water cycling at the sites, e.g. build up or leaching
- No changes in soil solution  $\text{NH}_4$  were observed in time although  $\text{NH}_4$  in deposition has declined

Vanguelova et al, 2010




## DON increase in soil solution - in response to higher DON input and or accumulation, recovery from acidity and increased microbial activity?





### Possible causes:

- 1) soil pollution recovery
- 2) increased microbial activity,
- 3) increased soil temperature
- 4) Increase in soil N accumulation


Vanguelova et al, 2010



## UK BioSoil plot locations - 167 plots



**Podzol**



**Gleysol**



**Histosol**







## Some conclusions

- Concerns over the local effects of excess nitrogen deposition and its forms remain and need to be taken into account in forest design plans around intensive farming in order to help N abatement but also to reduce the impact on the environment (forest, soils and waters).
- The importance of biological generated N should not be overlooked when considering N deposition impacts on soil processes and forest C balance.
- Long term trend analysis suggest significant increase in DON in rainfall, throughfall and soil solution in most of the UK Level II forest sites. The source of this organic dissolved nitrogen in rainfall is uncertain.
- Some preliminary results suggest that UK forest soils have accumulated N during the past 15-20 years and the likelihood for this N to be released are yet to be determined

- Forest Management Practices can override or mitigate these responses!!!



**BioSoil:**

Malcolm Reeves,  
Frank Heaven,  
Richard Hartnup,  
Andrew Hipkin  
Gordon Hudson  
Tanja Sanders

**Thanks to:****Intensive Monitoring**

Sue Benham  
Andy Moffat  
Dave Durrant  
Peter Crow  
Paul Taylor

**References:**

- Vanguelova, E.I., Nisbet, T.R., Moffat, A.J., Broadmeadow, S., Sanders, T.G.M. and Morison, J.I.L. (2013). A new evaluation of carbon stocks in British forest soils. *Soil Use and Management*, 29, 2, 169-181.
- Benham, S., Vanguelova, E. and Pitman, R. (2012). Short and long term changes in carbon, nitrogen and acidity in the mineral forest soils under oak at the Alice Holt Environmental Change Network site. *Science of the Total Environment* 421-422, 82-93.
- Vanguelova, E.I., Reynolds, B., Nisbet, T. and Godbold, D. (2011). The cycling of pollutants in non-urban forested environments. In: Levia, D.F., Carlyle-Moses, D.E. and Tanaka, T. (Eds.), *Forest Hydrology and Biogeochemistry: Synthesis of Past Research and Future Directions*. Ecological Studies Series, No. 216, Springer-Verlag, Heidelberg, Germany: DOI 10.1007/978-94-007-1363-5\_34.
- Vanguelova, E. and Pitman, R. (2011). Impacts of N inputs on forests and forest soil biogeochemistry in Great Britain. Paper in proceedings of "Nitrogen & Global Change – Key findings – future challenges", April 11-15, 2011, Edinburgh, UK.
- Vanguelova, E.I., Benham, S., Pitman, R., Moffat, A., Broadmeadow, M., Nisbet, T., Durrant, D., Barsoum, N., Wilkinson, M., Bochereau, F., Broadmeadow, S., Hutchings, T., Crow, P., Durrant-Huston, T., Taylor, P. (2010). Chemical fluxes in time through forest ecosystems in the UK – soil response to pollution recovery. *Environmental Pollution* 158 (5): 1857-1869.
- Vanguelova, E., Pitman, R., Luiro J. and Helmisaari H-S. (2010). Long term impacts of whole tree harvesting on soil carbon and nutrient sustainability in the UK. *Biogeochemistry* 101, 43-59.
- Pitman, R., Vanguelova, E.I. and Benham, S. (2010). Effects of phytophagous insects on the nutrient concentrations and fluxes through forest stands in the UK Level II network. *Science of the Total Environment* 409 (1), 169-181.
- Langan, S., Fransson, L. and Vanguelova, E. (2009). Dynamic modelling of UK forest soils to changes in acid deposition using the SAFE model. *Science of the Total Environment* 407, 5605-5619.
- Vanguelova, E.I., Barsoum, N., Benham, S., Broadmeadow, M., Moffat, A., Nisbet, T. and Pitman, R. (2007). Ten years of intensive environmental monitoring of British forests. Forestry Commission Information Note 88.